200400096

THE UNIVERD STAYLES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

Idula Research Joundation, Inc. (representing the interests of the Idula Agricultural Experiment Station and the University of Idula)

MICROS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FOM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR CONDITIONING IT OPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN PRODUCING A HYBRID OR T VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. IN STATES SEED OF THIS VARIETY (I) SHALL BE SOLD BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED SEED.

ALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER OF THE RIGHTS. (84 STAT.)

WHEAT, COMMON

'Idaho 587'

In Testimone Therest, I have hereunto set my hand and caused the seal of the Hant Harieto Protection Office to be affixed at the City of Washington, D.C. this twenty-third day of July, in the year two thousand and four.

Atlost:

Commissioner Plant Variety Protection Office Agricultural Marketing Service Agriculturo

2

2004 00 096

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 for issuance of the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority and provide evidence that name has been cleared by the appropriate recognized authority before the Certificate of Protection is issued. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, 10301 Baltimore Avenue, Suite 401 NAL Building, Beltsville, MD 20705. Telephone: (301) 504-5682 http://www.ams.usda.gov/lsg/seed.htm.

ITEM

- 19a. Give:
- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability, and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 19b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 19c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 19d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 19e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 20. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 23. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 24. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 22. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 23. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

24. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

US Patent 5,369,022. Filed September 17, 1993. Issued November 29, 1994. Assignee: BASF

Agrochemical Products, B.V. (Formerly American Cyanamid)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. The fees for filing a change of address; owner's representative; ownership or assignment; or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number for this information collection is 0581-055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

PVP Application No. 200400096 – Revised 29 April, 2004 Plant Variety Protection Application Exhibit A: Origin and Breeding History of the Variety - Idaho 587

Idaho 587 was derived from the third backcross of Stephens to the imidazolinone-resistant line 'FS4' (synonym Cv. 9804), with the backross having the pedigree Stephens*4/FS4. The imidazolinone resistance in FS4 was derived through mutagenesis of the imidazolinone-susceptible cultivar 'Fidel', followed by selection of the mutagenized population with imidazolinone herbicides (Newhouse 1992). The herbicide tolerance carried in 'FS4' is conditioned by an allele of the Als1 locus for acetolactate synthase with the allele designation Als1fid04. The BC₃F₂ seed of the cross Stephens*4/ FS4 was developed under contract by the American Cyanamid Corporation (subsequently acquired by BASF) at Texas A & M University by Mark Lazar and transferred to the University of Idaho under a material transfer agreement for utilization in selection and breeding. In 1996, the BC₃F₂ population designated 9697WAG460 was planted at Aberdeen, Idaho, and individual plants similar in appearance to Stephens were selected in fall 1997. BC₃F₃ seed of individual plant selections were planted at Aberdeen in fall 1997, with a random selection of heads collected in fall 1998 and planted as headrows at Aberdeen in fall 1998. Twenty BC₃F₃ seed of individual plant selections were planted in the greenhouse in winter 1999 and treated with imazamox (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-(methoxymethyl)-3-pyridinecarboxylic acid, BASF Corp) to identify BC₃F₂ families homozygous for Als1fid04. Headrows derived from BC₃F₂ families homozygous for Als1fid04 were selected in spring 1999 for uniformity and general agronomic appearance. Among these rows, a row designated W99-12-6 was harvested in 1999 and placed into yield testing and herbicide tolerance trials in southern Idaho in fall 1999. W99-12-6 was yield tested Aberdeen in 1999-2000, then placed in multi-location yield trials in 2000-2001. In 2001 and 2002, W99-12-6 was evaluated under the breeding line number IDO587 in cooperative testing with Oregon State University, University of Idaho Extension Testing, and the University of Idaho Weed Science Program. In fall 2000, 400 heads were selected from W99-12-6 and planted in the field at Aberdeen that same year. In spring 2001, the headrows were selected with imazamox herbicide for uniformity of tolerance. Headrows were uniformly tolerant to imazamox. Any headrows not similar in appearance to Stephens were discarded. Remaining BC₃F_{3:6} headrows were harvested and planted at Aberdeen in fall 2001 to form breeder seed of Idaho 587. Idaho 587 has been stable and uniform for herbicide tolerance and appearance for four years, without variants.

Plant Variety Protection Application Exhibit B: Statement of Distinctness Idaho 587

Idaho 587 is most similar to the cultivar 'Stephens'. It is a backcross derivative of Stephens that differs from the recurrent parent in only minor phenotypic variation. Idaho 587 is distinct from Stephens in being homozygous for the *Als1fid04* allele conditioning tolerance to the herbicide imazamox (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-(methoxymethyl)-3-pyridinecarboxylic acid, BASF Corp). Stephens is homozygous for an alternate allele that is normally considered a wild-type allele conditioning susceptibility to the imazamox herbicide (See Tables 8 and 9 in Exhibit D).

Form Approved - OMB No. 0581-005

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this collection of information is (0581-0055). The time required to complete this information collection is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marrial or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705

EXHIBIT C (Wheat)

OBJECTIVE DESCRIPTION OF VARIETY

		WHEAT	(Triticum spp.)		
NAME OF APPLICANT(S)				FOR OFFICIAL USE ONLY	
Id	daho Agricultura	l Experiment	t Station	PVPO NUMBER	
ADDRESS (Street and No. or RD	No., City, State, and Zip Code)			2004 0	0 096
	D Box 442337 DSCOW ID 83844-2	337		VARIETY NAME	
	٠			TEMPORARY OR EXPERIMEN	ral designation
a minimum of 100 plants. may be used to determine	STRUCTIONS CAREFULLY: (e.g. 0 9 9 or 0 9) wh Comparative data should be dete plant colors; designate system use is for your variety; lack of response	en number is either 99 o rmined from varieties er d:	r less or 9 or less respectively. stered in the same trial. Royal	Data for quantitative plant cha	racters should be based on
1. KIND:					
1	1=Common	2=Durum	3=Club	4=Other	(SPECIFY):
2. VERNALIZATI	ON:				
2	1=Spring	2=Winter	3=Other (S	SPECIFY):	· · · · · · · · · · · · · · · · · · ·
3. COLEOPTILE A	ANTHOCYANIN:				
1	1=Absent	2=Present			
4. JUVENILE PLA	NT GROWTH:				
1	1=Prostrate	2=Semi-erect	3=Erect		
5. PLANT COLOR	(boot stage):	· · · · · · · · · · · · · · · · · · ·			
2	1 = Yellow-Green	2 = Green	3 = Blue-Gr	reen	
6. FLAG LEAF (bo	oot stage):	:			
1	1 = Erect	2 = Recurved	2	1 = Not Twisted	2 = Twisted
7. EAR EMERGEN	CE:				
0 2	Number of Days Earl	ier Than	Stephens		*
0 3	Number of Days Late	r Than	Brundage		*

S&T-470-6 (2-99) designed by the Plant Variety Protection Office with WordPerfect 6.0a. Replaces LMGS-470-6 (6-82) which is obsolete

					. ***		49
8.	ANTHER COLOR:						
	1 =	Yellow	2 = Purple				
9.	PLANT HEIGHT (from	soil to top of hea	d, excluding awns)	:			···
	0 4 cm	Taller Than	Brundage 9	6		· · · · · · · · · · · · · · · · · · ·	*
	1 0 cm	Shorter Than _	Hubbard		· · · · · · · · · · · · · · · · · · ·		*
				* Relative to	a PVPO-Approved (Commercial Variety Gro	wn in the Same Tri
10.	STEM:		<u> </u>		· · · · · · · · · · · · · · · · · · ·		
	A. ANTHOCYAN	IN .		D. INTERNO	DE (SPECIFY	NUMBER)	
	1 I= Absent	2=Present		1 I= Hol	low 2=Se	mi-solid	3=Solid
	B. WAXY BLOOM	1		E. PEDUNCL	E		
	1=Absent	2=Present		1 1=Abse	ent 2=Pr	esent	
	C. HAIRINESS (la	ast internode of ra	ichis)	13 cm Len	igth		
	1=Absent	2=Present	• .				
11.	HEAD (at Maturity):						
	A. DENSITY			C. CURVATU	RE		
	2 1=Lax 3= Dense	2=Middense	•	$\boxed{2}$ 1 = Ere	eet 2 = In	ıclined	3 = Recurved
	B. SHAPE			D. AWNEDNE	ESS		
	$ \begin{array}{ c c } \hline 2 & 1 = \text{Tapering} \\ 3 = \text{Clavate} \end{array} $	g 2= Strap 4 = Other (S	PECIFY):	$ \begin{array}{ c c } \hline 4 & 1 = Aw \\ 3 = Aw \end{array} $		pically Awnletted	
·							
12.	GLUMES (at Maturity)	:					
	A. COLOR			C. BEAK			
	1 = White	2 = Tan		2	1 = Obtuse 3 = Acuminate	2 = Acute	
	3 = Other (S	PECIFY) :			·		
	B. SHOULDER			D. LENGTH			
	1 = Wanting 3 = Rounded 5 = Elevated	4 = Square		2	1 = Short (ca. 7mm) 3 = Long (ca.		
				,			

	·	~~~					<u> </u>	····
12.		S (at Maturity)	Continued:				2004 00	096
	E. W	IDTH					- ·	
	2	1 = Narrow (3 = Wide (ca		Medium (ca. 3.5m)	m)	•		
13.	SEED:						>	
	A. SI	HAPE			C. BI	RUSH		
	1	1 = Ovate	2 = Oval	3 = Elliptical	2	1=Short	2=Medium	3=Long
					1	1 = Not Collared	2 = Collared	
	B. CF	HEEK			D. CF	REASE		
	1	1=Rounded	2=Angu <u>l</u> ar		2	1 = Width 60% or le 2 = Width 80% or le 3 = Width Nearly as	ss of Kernel	• • • • • • • • • • • • • • • • • • • •
					1	1 = Depth 20% or les 2 = Depth 35% or les 3 = Depth 50% or les	s of Kernel	
	E. Col	or			G. PH	ENOL REACTION (see instructions):	
	1	1=White, 4= OTHER (S	2= Amber Specify)	3= Red	3	1 = Ivory 3 = Light Brown 5 = Black	2 = Fawn 4 = Dark Brown	1
*	F. TE	XTURE		•				
	2	1=Hard	2=Soft					
14. D	ISEASE:	(0=Not To	ested; 1=Susce	ptible; 2=Resis	tant; 3	=Intermediate; 4=1	Folerant)	
			PLEASE INDIC	ATE THE SPECI	FIC RAC	CE OR STRAIN TEST	ED	
	0	Stem Rust (Pi	uccinia graminis (2	Leaf Rust (Puccinia		
	2	Stripe Rust (I	Puccinia striiform	is)	0	Loose Smut (Ustilage	o tritici)	
	ď	Tan Spot (Pyr	renophora tritici-r	epentis)	0	Flag Smut (Urocystis	agropyri)	
	0	Halo Spot (Se	lenophoma donac	cis)	2	Common Bunt (Title	tia tritici or T. laevis,)
	1	Septoria nodor	rum (Glume Blot	ch)	1	Dwarf Bunt (Tilletia	controversa)	
	0	Septoria avena	e (Speckled Leaf	Disease)	0	Karnal Bunt (Tilletia	indica)	
	. 1	Septoria tritici	(Speckled Leaf I	Blotch)	0	Powdery Mildew (Er	y <i>siphe graminis</i> f. sp	. tritici)

"Snow Molds"

Scab (Fusarium spp.)

0

						2004 00 096
14.	Disea	se (Continued)	(0=Not Tested; 1=Su	sceptible; 2=	Resistant;	3=Intermediate; 4=Tolerant)
		·	PLEASE INDICA	TE THE SPECI	FIC RACE	OR STRAIN TESTED
	3	"Black Point"	(Kernel Smudge)	3	Common Bipolaris	Root Rot (Fusarium, Cochliobolus and spp.)
	0	Barley Yellow	Dwarf Virus (BYDV)	0	Rhizoctor	nia Root Rot (Rhizoctonia solani)
	0	Soilborne Mosa	ic Virus (SBMV)	4	Black Cha	aff (Xanthomonas campestris pv. translucens)
	0	Wheat Yellow	Spindle Streak) Mosaic Vi	rus 0	Bacterial syringae)	Leaf Blight (Pseudomonas syringae pv.
	0	Wheat Streak I	Mosaic Virus (WSMV)		Other (Sl	PECIFY)
च्यापः । ^र		Other (SPECII	FY)		Other (SI	PECIFY)
		Other (SPECIA	?Y)		Other (SI	PECIFY)
		Other (SPECIA	FY)		Other (SI	PECIFY)
15.	INSECT:	(0=Not Test	ed; 1=Susceptible; 2=	=Resistant; 3=	=Intermedi:	ate; 4=Tolerant)
•			PLEASE SPEC	Г РУ ВІОТУРЕ (where need	ed)
1.	0	Hessian Fly (M	ayetiola destructor)		Other (SI	PECIFY)
. • .	0	Stem Sawfly (C	ephus spp.)		Other (SP	PECIFY)
	0	Cereal Leaf Bee	tle (Oulema melanopa)		Other (SP	ECIFY)
	0	Russian Aphid	(Diuraphis noxia		Other (SP	ECIFY)
	0	Greenbug (Schi	zaphis graminum)		Other (SP	ECIFY)
	0	Aphids	· · · · · · · · · · · · · · · · · · ·		Other (SP	ECIFY)

Plant Variety Protection Application Exhibit D: Additional description Idaho 587

- Table 1. Group 5 (Imidazlinone Trial), Aberdeen, untreated controls, 2001-2.
- Table 2. Fidel, IDO587, and IDO588 treated with Imazamox herbicide at four rates: ANOVA, 2002.
- Table 3. Injury and agronomic performance for genotypes treated with differential rates of imazamox, Aberdeen, 2002.
- Table 4a. Mean Effect of Rate on Fidel, IDO587, and IDO588 (averaged over genotype), Aberdeen, 2002.
- Table 4b. Effect of Imazamox Rate on Stephens, Aberdeen, 2002.
- Table 5. Analysis of Variance, Mixed Effects Model, Aberdeen, 2000-01.
- Table 6. Interaction means (genotype *rate) for injury 22 DAT and heading date, Aberdeen, ID, 2000-1.
- Table 7. Grain yield of Stephens-derived breeding lines with imazamox resistance, tested with varying rates of imazamox herbicide, Aberdeen, ID, 2000-1.
- Table 8. Analysis of variance, mixed effects model, for IDO587, IDO588, and Fidel Fs-4, Pendelton, OR, 2001-2
- Table 9. Grain yield of IDO587 and IDO588 breeding lines with imazamox resistance, tested with varying rates of spring applied imazamox herbicide, Pendelton, OR, 2001-2
- Table 10. Western Regional Soft White Winter Wheats & Name Varieties, 2 YR SUMMARY
- Table 11. Milling and baking data from eight southern Idaho locations of extension nurseries, 2002.
- Table 12. Summary of north Idaho extension testing, Nez Perce, Lewiston, Genesee, and Bonner's Ferry, ID, 2001-2.
- Table 13. Irrigated winter wheat trials, Aberdeen and Hazelton, 2001 and 2002.
- Table 14. Summary of yield trial data from breeding program trials in Idaho in 2001 and 2002 and extension trials in Idaho, 2002.

Table 1. Group 5 (Imidazlinone Trial), Aberdeen, untreated controls, 2001-2.

	Head date	Yield	Tare	Test Weight
	(Julian)	(bu/a)		(lb/bu)
Fidel	163	121	3.6%	59.8
Stephens	165	121	5.2%	58.7
IDO587	165	132	3.1%	59.2
IDO588	164	123	1.9%	59.1
LSD Stderr	2	13	2%	
Stuerr	•			0.2

Table 2. Fidel, IDO587, and IDO588 ANOVA and Interaction Means, 2002.

Height Genotype (June 5) 2 174.0 87.0 2.68 (June 5) Rate Rate*Genotype 3 1583.5 527.8 29.67° Rate*Genotype 6 207.6 34.6 1.94 Heading Genotype 2 17.0 8.5 4.05 (Julian) Rate 3 21.2 7.1 3.94° Rate*Genotype 6 4.0 0.7 0.37 Injury Genotype 2 0.2 0.1 0.25 (May 24) Rate 3 26.4 8.8 19.49* Rate*Genotype 6 0.7 0.1 0.25 Injury Genotype 2 0.0 0.0 0.09 (May 31) Rate 3 57.0 19.0 22.43*	
Rate*Genotype 6 207.6 34.6 1.94 Heading Genotype 2 17.0 8.5 4.05 (Julian) Rate 3 21.2 7.1 3.94* Rate*Genotype 6 4.0 0.7 0.37 Injury Genotype 2 0.2 0.1 0.25 (May 24) Rate 3 26.4 8.8 19.49* Rate*Genotype 6 0.7 0.1 0.25 Injury Genotype 2 0.0 0.0 0.09	8
Heading Genotype 2 17.0 8.5 4.05 (Julian) Rate 3 21.2 7.1 3.94 Rate*Genotype 6 4.0 0.7 0.37 (May 24) Rate 3 26.4 8.8 19.49 Rate*Genotype 6 0.7 0.1 0.25 Rate*Genotype 6 0.7 0.1 0.25 (Injury Genotype 2 0.0 0.0 0.09	***
(Julian) Rate Rate*Genotype 3 21.2 7.1 3.94* Rate*Genotype Injury Genotype 2 0.2 0.1 0.25 Rate 3 26.4 8.8 19.49* Rate*Genotype 6 0.7 0.1 0.25 Injury Genotype 2 0.0 0.0 0.09	4
(Julian) Rate Rate*Genotype 3 21.2 7.1 3.94* Rate*Genotype Injury Genotype 2 0.2 0.1 0.25 Rate 3 26.4 8.8 19.49* Rate*Genotype 6 0.7 0.1 0.25 Injury Genotype 2 0.0 0.0 0.09	5
Rate*Genotype 6 4.0 0.7 0.37 Injury Genotype 2 0.2 0.1 0.25 (May 24) Rate 3 26.4 8.8 19.49* Rate*Genotype 6 0.7 0.1 0.25 Injury Genotype 2 0.0 0.0 0.09	
Injury Genotype 2 0.2 0.1 0.25 (May 24) Rate 3 26.4 8.8 19.49* Rate*Genotype 6 0.7 0.1 0.25 Injury Genotype 2 0.0 0.0 0.09	
(May 24) Rate 3 26.4 8.8 19.49* Rate*Genotype 6 0.7 0.1 0.25 Injury Genotype 2 0.0 0.0 0.09	ŗ
Rate*Genotype 6 0.7 0.1 0.25 Injury Genotype 2 0.0 0.0 0.09	5
Rate*Genotype 6 0.7 0.1 0.25 Injury Genotype 2 0.0 0.0 0.09	***
(14 04) = 1	
(14.00)	1
22,40	
Rate*Genotype 6 1.1 0.2 0.22	1
Injury Genotype 2 0.8 0.4 3.35	5
(June 10) Rate 3 142.9 47.6 63.31**	***
Rate*Genotype 6 4.5 0.8 1.01	
Injury Genotype 2 0.4 0.2 0.77	,
(June 17) Rate 3 177.7 59.2 167.27*	
Rate*Genotype 6 3.5 0.6 1.63	
7.03	,
Yield Genotype 2 750.0 375.0 1.92	! -
Rate 3 7145.0 2381.7 27.79**	***
Rate*Genotype 6 176.0 29.3 0.34	
Test Wt Genotype 2 1.94 0.97 1.74	
Rate 3 52.40 17.47 59.89**	
Rate*Genotype 6 5.00 0.83 2.85*	

Page 1 of

Genotype	Rate	Height June 5	Heading date	Injury May 24	Injury May 31	Injury June 10	Injury June 17	Viely	Test
		(cm)	(Julian)	(6-0)	(6-0)	(6-0)	(6-0)	(b) (d)	
Fidel	0	69.3	163	0.0) O) c	0 0	124.0)	
Fidel	0.04	67.0	164	0.5	0.0	9 0	2 0	7 1.0	0.0
Fidel	0.078	61.3	163		, ,	σ	. t	5.5.5	0.00
Fidel	0.156	47.3	162	2.0	2.5	. 4 5 6	- 4 . r.	0.20	00.0 76.0
IDO587	0	0.09	165	0.0	00	2 0	2 6	25. 5	20 0
IDO587	0.04	60.5	165	. C			9 6	5 6	20.0
IDO587	0.078	56.3	165	, ,	, ,	5 4) ; ;	55.9	200 200 200 200 200 200 200 200 200 200
IDO587	0.156	200	763	- -	- ر ن ر	D (7.0	114./	57.9
00000	3 ,	0	3	œ.	7.5	3.8	4. 8.	101.6	26.7
222	0	61.3	164	0.0	0.0	0.0	0.0	122.8	59.1
DO588	0.04	8.09	164	0.3	0.0	0.3	0.3	130.5	59.1
DO588	0.078	59.5	163	1.0	0.8	0.5	& C	109.7	200
DO588	0.156	50.0	163	2.0	3.0	4.8	5.0	98.5	57.2
StdErr		2.4	₹	ç		,			

ဖ
φ
S
æ
ğ
ф.

Table 4a.	Table 4a. Mean Effect of Rate on Fidel, IDO587, and IDO588 (averaged over genotype), Aberdeen, 2002	Ite on Fidel,	IDO587, and	d IDO588 (a	averaged ov	er genotyp	e). Aberde	en. 200	2
		Height	Heading	Injury	Injury	Injury	Injury	-	Test
	Rate	June 5	date	May 24	May 31	June 10	June 17	Yield	weight
		(cm)	(Julian)	(6-0)	(6-0)	(6-0)	(6-0)	(bu/a)	(bu/a) (lb/bu)
	0	63.5	164	0.0	0.0	0.0	0.0	125	59.4
	0.04	62.8	164	0.4	0.0	0.1	0.1	130	59.2
	0.078	59.0	163	1.3	1.0	1.	4.	109	58.4
	0.156	49.1	163	1.9	2.7	4.3	8.4	6 5	56.7
	GS7	3.5	-	9.0	0.8	0.7	0.5	8	
	StdErr	٠			٠				0.2
Table 4b.	Table 4b. Effect of Imazamox Rate on Stephens, Aberdeen, 2002	x Rate on S	stephens, Ab	erdeen, 200	25.	٠			
		Height	Heading	Injury	Injury	Injury	Injury		Test
	Rate	June 5	date	May 24	May 31	June 10	June 17	Yield	weight
		(cm)	(Julian)	(6-0)	(6-0)	(6-0)	(6-0)	(bu/a)	(nq/ql)
	0	56.8	165	0	0	0.0	0.0	121	
	0.04	23.3		4	6.5	7.0	8.0		•
	0.078	24.5		വ	6.5	7.3	8.5		
	0.156	21.5		9	7.5	8.0	8.8	•	
	US.)	4			, c				

Table 5. Analysis of Variance, Mixed Effects Model, Aberdeen, 2000-01

	Heading date		7 70	L/5 US	*** 60 75		0.8/ US/
	Height 31 DAT		ı	2	**		2
	Height	ш	900	0.0	34.67	0	0.0
	Injury 31 DAT		2		**	ú	
-	Injury 3	ட	0.56	9	126.04	77	-
2000-0	2 DAT	٠	SU			Š	
מוממנו, י	Injury 22 DAT	Ц	0.72	1 1	187.58	1.08	
ָבָּ בַּלָּנָלָ	8 DAT		us	**		SU.	
mand Theory Indeed Conference (And Conference)	Injury 16 DAT	T	1.07	470 CE	17.03	0.53	
(1)	njury 8 DAT		SI	***		SU	
	Injury 8	<u>ا</u>	1.03	21.04	P	0.85	
	+2691	בוופנו	Genotype	Rate	*	Genotype Rate	

Split plot design. Main plots = genotypes (Fidel and six Imi-Stephens selections) Subplots = rates (0, 5, 10, 20 oz product/a with 2.5% UAN and 0.25% NIS) 4 replications

Table 6. Interaction means (genotype *rate) for injury 22 DAT and heading date, Aberdeen, ID, 2000-1.

1		Injury	njury 22 DAT (0 to 9)	to 9))		Heading date (I	(acil	
		Rate	(OZ prod)	(int/a)					Julian	
1 1 1 1 1 1 1	<		10 P				Kate	oz prod	uct/a)	
genotype	٥	5	10	20	Mean	0	ιc	5	۶	Moon
Fidel	0.5	0.5	2.0	4.0	200	162	163	163	405	100
10.40.0	0	c	•		2	7	3	3	00	103
7-71 2	0.0	χ. Ο	 	4.3	9.	163	164	165	169	787
IS 12-4	0.0	0.8	20	4.8	0	30,	. 0	9 6	3 !	3 :
100) (į) F		60	<u>0</u>	108	1/1	99
) (2007)	0.0	0.5	<u>_</u> ∞	4.5	1.7	164	165	167	160	98
10.10,11	0	4	L	(,)	2	2	2	00
7 2 2	5	0.0	<u>.</u>	3.5	4.	165	165	167	168	166
5 12-13	<u>_</u>	٥,	ά	11	,	,			2	2
. () i	2	<u>,</u>	t,	\' .	164	164	166	169	166
DO588	0.0	80	23	eri eri	<u>ب</u>	185	707	700	1) (
				3	2	3	104	80	2	<u>/</u> 9
Mean	0.1	9.0	8.	4.1		164	16,	466	6	
			! •			5	5	20	80	

Table 7. Grain yield of Stephens-derived breeding lines with imazamox resistance, tested with varying rates of imazamox herbicide, Aberdeen, ID, 2000-1.

Rate lbs ai/ac	Fidel	IS1202	IS1204	IDO587	IS1211	IS1213	IDO588
				bu/ac			
0	93.5	111.3	116.4	126.3	113.0	116.1	117.9
0.04	96.2	107.9	109.0	117.1	106.6	116.3	109.8
0.078	86.3	110.9	115.6	115.1	115.2	107.6	104.5
0.156	86.8	91.5	101.1	99.3	103.8	117.2	107.0






Table 8. Analysis of variance, mixed effects model, for IDO587, IDO588, and Fidel Fs-4, Pendelton, OR, 2001-2

	Degrees	Finalisa	Drob > F			
	of freedom	F-value	Prob > F			
Rate	3	5.88	0.0025			
Genotype	2	6.41	0.0045			
Rate*Genotype	6	1.31	0.2815			
Data provided by Jim Peterson, Oregon State University						

Table 9. Grain yield of IDO587 and IDO588 breeding lines with imazamox resistance, tested with varying rates of spring applied imazamox herbicide, Pendelton, OR, 2001-2

Rate				
4 oz ai/ac	Fidel	IDO587	IDO588	Stephens
		b	u/ac	. "
0	82.7	78.9	83.7	77.6
4	89.8	82.7	93.4	0.0
6	90.4	79.2	82.7	0.0
12	85.7	71.8	72.4	0.0

Yield of 12 oz/ac trea	atment as %	% of untreated ch	eck.	
	104%	91%	86%	0%

Table 10. Western Regional Soft White Winter Wheats & Name Varieties, 2 YR SUMMARY Aberdeen and Hazelton, ID

	Solvent retention capacity Flour yield		yield	ı				
Genotype	Water	NaCarb	Sucrose	Lactic acid	Break	Total	Flour protein	Sugar snap cookie dia
				%				cm
Year	CONTROL OF THE PROPERTY OF THE	errelena v Arbeitela Arbeitela (1885)	WO 45 WINDOWS TO THE TOTAL PROPERTY OF THE T	er vonum vondamning Animinal AAA vidal (NAAA)	M MO STORES NATIONAL SECTION CONTRACTOR CONT	or or occurrence and other life	AMA-ALMATAN'A YARUTUNYA TITOTA TOTOTA	
2001	49.9	62.1	93.6	84.2	34.3	63.0	10.8	8.2
2002	48.9	65.9	97.1	87.9	39.9	67.1	9.7	8.4
Location								
Aberdeen	48.1	63.3	91.7	81.6	36.5	66.3	10.1	8.3
Hazelton	50.6	64.7	98.9	90.5	37.7	63.7	10.4	8.2
Genotype								and it for the form
Brundage	49.0	63.4	91.8	85.8	38.0	65.7	9.9	8.4
Brundage96	47.1	62.6	93.7	87.5	36.8	63.2	10.6	8.3
Daws	50.2	64.6	102.3	98.4	39.1	63.0	9.8	8.1
Madsen	51.4	62.9	93.0	79.4	35.9	66.5	10.5	8.4
Stephens	51.6	66.1	97.1	94.0	33.8	63.8	10.6	8.1
Weatherford	51.6	64.5	96.8	85.6	34.3	66.0	10.5	8.2
WPB470	47.7	66.4	100.3	101.9	35.4	63.1	10.8	8.0
IDO587	48.5	64.3	92.7	80.0	34.1	65.5	10.0	8.5
IDO605	48.2	61.8	90.3	71.6	34.6	68.4	10.5	8.3

Table 11. Milling and baking data from eight southern Idaho locations of extension nurseries, 2002.

Cultivar	Break flour yield	Flour yield	Flour protein	Cookie diameter	Top grain
	%	%	%	cm	1 to 5
Idaho 587	35.9	65.9	10.3	8.5	2.7
Stephens	35.9	65.2	10.2	8.4	2.8
Brundage	38.1	64.8	9.9	8.4	2.4
Std error	0.6	0.6	0.2	0.9	0.3

Table 12. Summary of north Idaho extension testing, Nez Perce, Lewiston, Genesee, and Bonner's Ferry, ID, 2001-2.

Cultivar	Grain yield bu/ac	Test weight lbs/bu	Grain protein %	Lodging ¹ %	Plant height
Stephens	97	57.7	11.2		<u>in</u>
· ·				•	36
IDO587	96	57.8	11.3	1	36
IDO588	99	58.4	11.4	3	37
Albion	94	56.2	11.3	3	34
Brundage 96	99	58.1	11.3	0	35
Beamer	101	58.4	11.5	12	38
Finch	97	58.7	11.5	1	38
Hubbard	100	59.4	11.2	0	40
Lambert	100	58.3	11.3	5	39
Madsen	96	57.9	11.9	2	36
Mohler	109	58.3	10.9	6	38
Rod	98	57.6	10.7	3	35
Tubbs	102	56.4	11.7	0	37
WPB470	101	61.5	11.3	2	35
Std. Err.	2.1	0.3	0.2		0.5

1. Lodging data for Bonner's Ferry only.

Table 13. Irrigated winter wheat trials, Aberdeen and Hazelton, 2001 and 2002.

		Grain	Test			Lodging
	•	yield	weight	Heading	Height	1=erect
		bu/ac	#/bu	Jan 1+	in	9=Flat
Brundage	SWW	123.9	62.8	155.3	32.1	2.0
IDO605	SWW	118.8	60.0	163.1	35.6	2.2
IDO587	SWW	117.1	. 60.3	158.3	33.6	2.4
Stephens	SWW	114.2	59.7	160.7	35.7	3.5
Boundary	HRW	113.7	61.9	159.8	33.8	3.4
Garland	HRW	112.6	61.1	160.0	28.1	1.5
Moreland	HRW	111.1	61.3	155.1	32.0	2.2
WPB470	SWW	109.8	63.2	155.3	32.0	1.8
Brundage 96	SWW	101.3	59.4	161.8	32.5	3.2
Std. Error		12.3	0.5	2	2.0	

Table 14. Summary of yield trial data from breeding program trials in Idaho in 2001 and 2002 and extension trials in Idaho, 2002.

	Rain-fed sites #	Irrigated sites #	Grain yield bu/ac	Test weight lbs/bu	Plant height in	Lodging %	Heading date Jan 1+
IDO587	4	10	106.4	57.6	33.5	16	159
Stephens	4	10	107.1	57.8	34.1	21	160
			ns	ns	ns	ns	ns
IDO587	1	9	107.6	57.1	32.9	24	159
Brundage	1	9	113.5	60.0	31.7	18	156
,			ns	**	*	ns	**
IDO587	4	10	106.4	57.6	33.5	16	159
Brundage 96	4	10	101.5	57.8	32.7	16	161
_			ns	ns	*	ns	ns
IDO587	4	6	103.7	57.0	33.6	18	169
Hubbard	4	6	105.7	59.6	38.3	19	162
			ns	**	**	ns	**
IDO587	4	7	94.7	56.4	32.0	5	160
Madsen	4	7	93.6	57.4	33.0	0	164
			ns	ns	ns	ns	**
IDO587	4	10	106.4	57.6	33.5	16	159
WPB470	4	10	106.0	61.9	32.2	12	156
			ns	**	**	ns	**

ns= the difference between paired cultivars is not signficant at the 95% level of confidence

^{*=}difference between paired cultivars significant at the 95% level of confidence.

^{**=}difference beween paired cultivars significant at the 99% level of confidence.

IDAHO AGRICULTURAL EXPERIMENT STATION Moscow, Idaho

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

Announces the release of IDAHO 587 SOFT WHITE WINTER WHEAT

Idaho 587, soft white winter wheat (*Triticum aestivum*, L.), was developed by the Idaho Agricultural Experiment Station for use by grain producers in the Pacific Northwest. Idaho 587 is a backcross derivative of the soft white winter wheat cultivar 'Stephens' with genetic tolerance developed by the BASF Corporation to the imidazolinone class of herbicides for use in controlling pernicious weeds of winter wheat and generically given the name 'Clearfield'.

Idaho 587 was derived from the third backcross of Stephens to the imidazolinone-resistant line 'FS4' (synonym Cv. 9804), with the backross having the pedigree Stephens*4/FS4. The imidazolinone resistance in FS4 was derived through mutagenesis of the imidazolinone-susceptible cultivar 'Fidel', followed by selection of the mutagenized population with imidazolinone herbicides (Newhouse 1992). The herbicide tolerance carried in 'FS4' is conditioned by an allele of the Als1 locus for acetolactate synthase with the allele designation Als1fid04. The BC₃F₂ seed of the cross Stephens*4/ FS4 was developed under contract by the American Cyanamid Corporation (subsequently acquired by BASF) at Texas A & M University by Mark Lazar and transferred to the University of Idaho under a material transfer agreement for utilization in selection and breeding. In 1996, the BC₃F₂ population designated 9697WAG460 was planted at Aberdeen, Idaho, and individual plants similar in appearance to Stephens were selected in fall 1997. BC₃F₃ seed of individual plant selections were planted at Aberdeen in fall 1997, with a random selection of heads collected in fall 1998 and planted as headrows at Aberdeen in fall 1998. Twenty BC₃F₃ seed of individual plant selections were planted in the greenhouse in winter 1999 and treated with imazamox (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-(methoxymethyl)-3-pyridinecarboxylic acid, BASF Corp) to identify BC₃F₂ families homozygous for Als1fid04. Headrows derived from BC₃F₂ families homozygous

for Als1fid04 were selected in spring 1999 for uniformity and general agronomic appearance. Among these rows, a row designated W99-12-6 was harvested in 1999 and placed into yield testing and herbicide tolerance trials in southern Idaho in fall 1999. W99-12-6 was yield tested Aberdeen in 1999-2000, then placed in multi-location yield trials in 2000-2001. In 2001 and 2002, W99-12-6 was evaluated under the breeding line number IDO587 in cooperative testing with Oregon State University, University of Idaho Extension Testing, and the University of Idaho Weed Science Program. In fall 2000, 400 heads were selected from W99-12-6 and planted in the field at Aberdeen that same year. In spring 2001, the headrows were selected with imazamox herbicide for uniformity of tolerance. Headrows were uniformly tolerant to imazamox. Any headrows not similar in appearance to Stephens were discarded. Remaining BC₃F_{3:6} headrows were harvested and planted at Aberdeen in fall 2001 to form breeder seed of Idaho 587. Idaho 587 has been stable and uniform for herbicide tolerance and appearance for four years without variants.

Idaho 587 is most similar to the cultivar Stephens with an unpigmented coleoptile, dark green foliage, and a prostrate to semi-erect fall growth habit. Idaho 587 is a semi-dwarf, winter wheat that is approximately 85 cm tall at maturity, similar to Stephens and Madsen (both 86 cm), but taller than 'Brundage 96' (83 cm) and shorter than 'Hubbard' (97 cm). Like Stephens, Idaho 587 is a medium maturity cultivar heading 159 days after January 1st, approximately a day earlier than Stephens and four days earlier than 'Madsen'. Idaho 587 has broad, recurved flag leaves, is awned, with yellow anthers at anthesis and white colored chaff at maturity. Idaho 587 has large, plump, oval, soft white seed with a wide crease, short brush, mid-sized embryo, and an average seed size is 46 mg (compared with 42 mg for Stephens). The seed of Idaho 587 is similar to Stephens for polyphenol oxidase activity. In replicated evaluations in Moscow, Idaho, Lewiston, Idaho, Pullman, Washington, and Mount Vernon, Washington, Idaho 587 had both seedling and adult plant resistance to the dominant races of stripe rust [caused by *Puccinia striiformis* (Westend.), races PST-14, 22, 23, 26,

-58

35, 40, 41, 45, 53, 61, 74, 78, 85, 91, 92, 93, 97, 98, 99], without observable necrosis or pustule formation. Idaho 587's stripe rust resistance is similar to Stephens and likely derives from that cultivar. Other disease resistances and susceptibilities for Idaho 587 are expected to be similar to Stephens.

The primary difference between Idaho 587 and Stephens is Idaho 587's tolerance to imazamox herbicide. In two years of evaluation at Aberdeen, Idaho and one year of evaluation at Pendleton, Oregon and Moscow, Lewiston, and Nez Perce, Idaho, using one to four times the recommended rates of imazamox herbicide, Idaho 587 a did not have significantly different response to imazamox than FS4, the source of Idaho 587's herbicide tolerance. In two years of trials at Aberdeen, 45 g ai ha⁻¹ imazamox was sufficient to kill 100% of the Stephens check but did not significantly reduce grain yield of Idaho 587. At the same rate of herbicide, in a spring application at 6 site-years across Idaho and Oregon, Idaho 587 had an average grain yield of 6.1 Mg ha⁻¹ compared with 6.3 Mg ha⁻¹ for FS4 (not significantly different).

southeastern Idaho in 2001 and 2002, and in western and northern Idaho in 2002, Idaho 587, Stephens, Brundage 96, and 'Westbred 470' had average grain vields of 7.2 Mg ha⁻¹, 7.2 Mg ha⁻¹, 6.8 Mg ha⁻¹, and 7.1 Mg ha⁻¹, respectively. In the same trials Idaho 587, Stephens, Brundage 96, and Westbred 470 had average test weights of 741 kg m⁻³, 744 kg m⁻³, 743 kg m⁻³, and 796 kg m⁻³, respectively. These four cultivars were not found to differ significantly in their lodging resistance. In a subset of these Idaho trials (11 site-years), Idaho 587 had similar grain yields to Madsen, 6.4 Mg ha⁻¹ and 6.3 Mg ha⁻¹, respectively. Across nine locations, in Oregon extension testing, in 2002, Idaho 587, Stephens, and Madsen had average grain yields of 5.7 Mg ha⁻¹, 5.5 Mg ha⁻¹, and 5.8 Mg ha⁻¹, respectively. In two sites across two years of milling evaluations in Idaho, Idaho 587 had an average flour yield of 655 g kg⁻¹ of grain compared with 638 g kg⁻¹ for Stephens and 665 g kg⁻¹ for Madsen. Sugar snap cookie evaluations of the same flour for Idaho 587 produced an average cookie

In yield trials, without Imazamox applications (14 site-years), in

92	diameter of 8.5 cm compared with 8.1 cm for Stephens and 8.4 cm for Madsen.
93	Solvent retention capacity analysis of flour from Idaho 587 and Stephens
94	produced a sodium carbonate solvent retention of 643 g kg ⁻¹ for Idaho 587, 661 g
. 95	kg ⁻¹ for Stephens, and 629 g kg ⁻¹ for Madsen. Lactic acid retention in the same
96	analyses of 800 g kg ⁻¹ for Idaho 587, 940 g kg ⁻¹ for Stephens, 794 g kg ⁻¹ for
97	Madsen.
98	Foundation seed of Idaho 587 will be maintained by the Idaho Agricultural
99	Experiment Station and Plant Variety Protection sought for the cultivar. Idaho
100	587 carries a patented proprietary gene owned by BASF Corporation. Therefore,
101	it will be distributed only under a material transfer agreement in cooperation with
102	the BASF Corporation. Interested parties may contact the Idaho Research
103	Foundation, Moscow Idaho.
	Director, Idaho Agricultural Experiment Station Date
	Moscow Idaho

References

Newhouse, Keith E., Wendy A. Smith, Mark A. Starrett, Thomas J. Schaefer, and Bijay K. Singh (1992) Tolerance to Imidazolinone Herbicides in Wheat. Plant Physiology 100:882-886.

REPRODUCE LOCALLY, Include form number and edition date on all	reproductions.	ORM APPROVED - OMB No. 0581-0055	
U.S. DEPARTMENT OF AGRICULTURE			
AGRICULTURAL MARKETING SERVICE	Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). The information is held		
EXHIBIT E	confidential until the certificate is issu	ed (7 U.S.C. 2426).	
STATEMENT OF THE BASIS OF OWNERSHIP			
1. NAME OF APPLICANT(S) Idaho Research Foundation, Inc. (representing the Interests of the Idaho Agricultural Experiment Station and the University of Idaho.)	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME	
Agricultural Experiment Station and the University of Idaho.)	IDA0587	Idaho 587	
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (Include area code)	6. FAX (Include area code)	
PO Box 443447 University of Idaho	(208) 885-7173	(208) 885-6654	
Moscow, ID 83844-2337 USA	7. PVPO NUMBER 20	04 00 096	
8. Does the applicant own all rights to the variety? Mark an "X" in the	e appropriate block. If no, please expl a	in. YES NO	
Idaho 587 contains the patented gene <u>al</u> Filed September 17, 1993. Issued Novembroducts, B.V. (Formerly American Cyana	<u>lsl.</u> U.S. Patent 5,369 aber 29, 1994. Assigno	9,022.	
9. Is the applicant (individual or company) a U.S. national or a U.S. b	ased company? If no, give name of c	ountry. X YES NO	
10. Is the applicant the original owner?	NO If no, please answer one	of the following:	
a. If the original rights to variety were owned by individual(s), is (are) the original owner(s) a U.S. Nation	• •	
 b. If the original rights to variety were owned by a company(ies), 	· · · · · · · · · · · · · · · · · · ·		
YES	NO If no, give name of countr	у	
	_		
11. Additional explanation on ownership (Trace ownership from origin Original cross developed under contract			
Segregating population transferred to t		-	
transfer agreement. Selection and test	· .		
University of Idaho (Edward Souza) with	permission of BASF.	•	
	•		
	<u></u>		
PLEASE NOTE:			
Plant variety protection can only be afforded to the owners (not licens	ees) who meet the following criteria:		
 If the rights to the variety are owned by the original breeder, that penaltonal of a country which affords similar protection to nationals of 			
If the rights to the variety are owned by the company which employ nationals of a UPOV member country, or owned by nationals of a o genus and species.			
3. If the applicant is an owner who is not the original owner, both the o	original owner and the applicant must m	eet one of the above criteria.	
The original breeder/owner may be the individual or company who direct for definitions.	ected the final breeding. See Section 4	1(a)(2) of the Plant Variety Protection	
According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, a control number. The valid OMB control number for this information collection is 0591-0055, including the time for reviewing the instructions, searching existing data sources, gathering are	The time required to complete this information collect	tion is estimated to average 0.1 hour per response.	
The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and ac marital or family status, political beliefs, perental status, or protected genetic information. (No communication of program information (Braille, large print, audiotape, etc.) should contact US	t all prohibited bases apply to all programs.) Persor	ns with disabilities who require alternative means for	

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whilten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.

ST-470-E (04-03) designed by the Plant Variety Protection Office using Word 2000